

## Preliminary Exam May 2018

**Problem 1.** Let  $\sigma > 0$ . Let  $(f_k)_{k \in \mathbb{N}}$  be a sequence of functions  $f_k : \mathbb{R} \rightarrow \mathbb{R}$  with  $f_k(0) = 0$ . Moreover let  $(A_k)_{k \in \mathbb{N}} \subset [0, \infty)$  be a *bounded* sequence of real numbers such that

$$|f_k(x) - f_k(y)| \leq A_k |x - y|^\sigma \quad \text{for all } x, y \in \mathbb{R}.$$

- (a) Show that there exists  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that a subsequence  $f_{k_i}$  converges uniformly to  $f$  in every interval  $[-a, a]$ ,  $a > 0$ .  
 (b) Show that  $f$  satisfies

$$|f(x) - f(y)| \leq A |x - y|^\sigma$$

where  $A = \liminf_{k \rightarrow \infty} A_k$ .

**Problem 2.** Prove that if  $X$  is a metric space and  $f : X \times [0, 1] \rightarrow \mathbb{R}$  is a continuous function, then  $g : X \rightarrow \mathbb{R}$ , defined by  $g(x) = \sup_{t \in [0, 1]} f(x, t)$ , is continuous.

**Problem 3.** Prove (using only the material covered in the course) that there is no continuous and one-to-one function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ . **Hint:** Assume that such a function exists and then restrict the function to the unit circle in  $\mathbb{R}^2$ .

**Problem 4.** Suppose  $f : \mathbb{R}_+^2 \rightarrow \mathbb{R}$  is a continuous function defined on

$$\mathbb{R}_+^2 = \{(x, y) : x \in \mathbb{R}, y > 0\}.$$

Assume also that the limits

$$g(u, v) = \lim_{t \rightarrow 0} \frac{f((u+t)\cos v, (u+t)\sin v) - f(u\cos v, u\sin v)}{t},$$

and

$$h(u, v) = \lim_{t \rightarrow 0} \frac{f(u\cos(v+t), u\sin(v+t)) - f(u\cos v, u\sin v)}{t},$$

exist and define continuous functions  $g, h$  on the domain

$$D = \{(u, v) : u > 0, 0 < v < \pi\}.$$

Prove that the function  $f$  is differentiable on  $\mathbb{R}_+^2$ .

**Problem 5.** Let  $f \in C^2(\Omega) \cap C^0(\bar{\Omega})$ , where  $\Omega \subset \mathbb{R}^n$  is open and bounded. Let  $\Delta f = \sum_{i=1}^n \partial^2 f / \partial x_i^2$  be the Laplace operator.

- (a) Show that if for some  $\varepsilon > 0$  and  $x_0 \in \Omega$  we have  $\Delta f(x_0) \geq \varepsilon$ , then  $f$  has no local maximum at  $x_0$ .  
 (b) Conclude that if  $\Delta f(x) \geq \varepsilon$  for some  $\varepsilon > 0$  and all  $x \in \Omega$ , then we have  $\sup_\Omega f = \sup_{\partial\Omega} f$ .  
 (c) Conclude that if  $\Delta f(x) \geq 0$  for all  $x \in \Omega$ , then we have  $\sup_\Omega f = \sup_{\partial\Omega} f$ .

**Hint for part (c):** Observe that  $\Delta|x|^2 = 2n$ . Use it to modify a function  $f$  in (c) so that you can apply part (b).

**Problem 6.** For  $x = (x_1, x_2) \in \mathbb{R}^2$ , let  $|x| = \sqrt{x_1^2 + x_2^2}$ . Let  $D = \{x \in \mathbb{R}^2 : |x| < 1\}$  and let  $f : \bar{D} \rightarrow \mathbb{R}$  be continuous on  $\bar{D}$ . Prove that

$$\lim_{n \rightarrow \infty} \iint_D (n+2)|x|^n f(x) dA = \int_0^{2\pi} f(\cos t, \sin t) dt.$$